



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2123909 C 2002/10/01

(11)(21) **2 123 909**

(12) **BREVET CANADIEN  
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1992/12/07

(87) Date publication PCT/PCT Publication Date: 1993/06/24

(45) Date de délivrance/Issue Date: 2002/10/01

(85) Entrée phase nationale/National Entry: 1994/05/18

(86) N° demande PCT/PCT Application No.: EP 1992/002826

(87) N° publication PCT/PCT Publication No.: 1993/011773

(30) Priorité/Priority: 1991/12/18 (91311761.0) EP

(51) Cl.Int.<sup>5</sup>/Int.Cl.<sup>5</sup> A61K 31/57, A61K 31/165

(72) Inventeurs/Inventors:  
Carling, Christer Carl Gustav, SE;  
Trofast, Jan William, SE

(73) Propriétaire/Owner:  
Aktiebolaget Astra, SE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : NOUVELLE COMBINAISON DE FORMOTEROL ET DE BUDESONIDE

(54) Title: NEW COMBINATION OF FORMOTEROL AND BUDESONIDE

(57) Abrégé/Abstract:

Effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are used in combination for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 31/57 // (A61K 31/57 A61K 31:165)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/11773</b> <b>(43) International Publication Date:</b> 24 June 1993 (24.06.93) <b>2123909</b>
<b>(21) International Application Number:</b> PCT/EP92/02826 <b>(22) International Filing Date:</b> 7 December 1992 (07.12.92) <b>(30) Priority data:</b> 91311761.0 18 December 1991 (18.12.91) SE <b>(71) Applicant:</b> AKTIEBOLAGET ASTRA [SE/SE]; S-151 85 Södertälje (SE). <b>(72) Inventors:</b> CARLING, Christer, Carl, Gustav ; Backvägen 8, S-240 10 Dalby (SE). TROFAST, Jan, William ; Vapenkroken 34, S-226 47 Lund (SE). <b>(74) Agents:</b> HJERTMAN, Ivan et al.; AB Astra, Patent Department, S-151 85 Södertälje (SE).		<b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> NEW COMBINATION OF FORMOTEROL AND BUDESONIDE  <b>(57) Abstract</b>  Effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are used in combination for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.		

23940-804(S)

1

**New combination of formoterol and budesonide.**Field of the invention

This invention relates to improvements in the treatment of mild as well as severe asthma and other  
5 respiratory disorders. More particularly, it relates to the use of a bronchodilator in combination with a steroidal anti-inflammatory drug for the treatment of respiratory disorders such as asthma, and to pharmaceutical compositions containing the two active ingredients. It emphasizes the use of a long-  
10 acting bronchodilator which provides rapid relief of symptoms.

Background of the invention

There have recently been significant advances in our understanding of asthma. Despite many advances, both in awareness of the disease by doctors and patients alike, coupled  
15 with the introduction of very powerful and effective anti-asthma drugs, asthma remains a poorly understood and often poorly treated disease. Previously, contraction of airway smooth muscles has been regarded as the most important feature of asthma. Recently there has been a marked change in the way  
20 asthma is managed, stemming from the fact that asthma is recognized as a chronic inflammatory disease. Uncontrolled airway inflammation may lead to mucosal damage and structural changes giving irreversible narrowing of the airways and fibrosis of the lung tissue. Therapy should therefore be aimed  
25 at controlling symptoms so that normal life is possible and at the same time provide basis for treating the underlying inflammation.

The most common cause for poor control of asthma is poor compliance with the long-term management of chronic  
30 asthma, particularly with prophylactic treatments, such as inhaled steroids, which do not give immediate symptom relief.

23940-804(S)

2

Patients will readily take  $\beta_2$ -agonist inhalers, since these provide rapid relief of symptoms, but often do not take prophylactic therapy, such as inhaled steroids, regularly because there is no immediate symptomatic benefit. They also  
5 counteract down regulation of  $\beta_2$ -adrenoceptor agonists.

Formoterol, (N-[2-hydroxy-5-[1-hydroxy-2-[[2-(4-methoxyphenyl)-1-methylethyl]amino]ethyl]phenyl]formamide), is an adrenoceptor agonist which selectively stimulates  $\beta_2$ -receptors, thus producing relaxation of bronchial smooth  
10 muscle, inhibition of the release of endogenous spasmogens, inhibition of oedema caused by endogenous mediators, and increased mucociliary clearance. Inhaled formoterol fumarate acts rapidly, usually within minutes which gives the patient immediate confirmation that he has taken an adequate dose and  
15 thereby avoiding overdosing of both  $\beta_2$ -agonist and steroid. Inhaled formoterol also exerts a prolonged bronchodilation, which in clinical trials has been demonstrated as up to 12 hours.

Budesonide, (16,17-butyridenebis(oxy)-11,21-dihydroxypregna-1,4-diene-3,20-dione), may be given in a high  
20 inhaled dose (up to 2 mg daily) with very low systemic effects, possibly because of its rapid metabolism. The high rapid systemic elimination of budesonide is due to extensive and rapid hepatic metabolism. Long term clinical studies have  
25 shown that inhaled budesonide is a pharmacologically safe drug. High doses of inhaled budesonide are highly effective and well tolerated when used in oral steroid replacement therapy. Budesonide represents a logical safe and effective therapy for long term control of asthma.

30 The inhaled route of administration enables the dose to be delivered directly to the airways. By this type of administration, it is possible to give a small dose and thereby

23940-804(S)

3

minimizing unwanted side-effects. The drawbacks of the currently available bronchodilators are their relatively short duration of action. By using a compound with long duration e.g. formoterol it would be possible to avoid the nocturnal  
5 asthma, which so often causes considerable anxiety and debility to the patients. Formoterol gives less nocturnal waking than the commonly used short-acting agonists like salbutamol, terbutaline and the like. Formoterol has been registered for oral administration in Japan since 1986.

10               Pharmaceutical combinations of long-acting  $\beta_2$ -agonists and steroids are disclosed in two European applications, EP 416950 which discloses the combination of salmeterol and beclomethasone, and EP 416951 which discloses the combination of salmeterol and fluticasone propionate.

15               In Ann. Allergy 1989, 63 (3), p. 220-224 the use of a  $\beta_2$ -agonist, i.e. formoterol and a steroid, i.e. budesonide separately are mentioned. It is not disclosed a pharmaceutical combination including both formoterol and budesonide, or the use of the two compounds in combination therapy. The use of a  
20  $\beta_2$ -agonist and a steroid separately is also mentioned in Lung (1990), 168, no. supp, p. 105-110.

#### Outline of the Invention

The present invention is based on the concept of a novel combination therapy whereby formoterol (and/or a  
25 physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation. This combination has not only a greater efficiency and duration of bronchodilator action but the combination also has a rapid onset of action. This new  
30 feature is of utmost importance in order to establish a higher compliance for patients and it provides a rescue medicine

23940-804 (S)

4

thereby avoiding the necessity for the patient of carrying two different inhalers. This simplifies life for patients considerably and makes life more comfortable and secure. The rapid onset of the long-acting  $\beta_2$ -agonist gives the patient  
5 immediate confirmation that he has taken an adequate dose and thereby avoiding overdosing of both  $\beta_2$ -agonist and steroid. Since the use of formoterol instead of salmeterol gives a much more rapid onset, the combinations according to the invention have a number of advantages compared to the combinations  
10 disclosed in EP 416950 and EP 416951. The combination according to present invention permits a twice daily dosing regime as a basic treatment of asthma, particularly nocturnal asthma.

The present invention provides a medicament  
15 containing, separately, or together, (i) formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and (ii) budesonide for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

20 The invention also provides a pharmaceutical composition for administration by inhalation in the treatment of respiratory disorder which composition comprises formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.

25 According to another aspect of the invention there are provided pharmaceutical compositions comprising effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide as a combined preparation for simultaneous, sequential or separate  
30 administration by inhalation in the treatment of respiratory disorder.

23940-804 (S)

5

The invention further provides formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide for use in combination therapy by simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

Further the invention provides the use of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) in the manufacture of a medicament for combination therapy where formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder and the use of budesonide in the manufacture of a medicament for combination therapy where formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder.

The invention additionally relates to the use of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide in the manufacture of a medicament for combination therapy for simultaneous, sequential or separate administration of formoterol and budesonide by inhalation in the treatment of respiratory disorder.

According to a further feature of the invention there is provided a method of treating respiratory disorder which comprises the simultaneous, sequential or separate administration by inhalation of effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.

Suitable physiologically acceptable salts of formoterol include acid addition salts derived from inorganic and organic acids, such as the hydrochloride, hydrobromide,

23940-804 (S)

6

sulphate, phosphate, maleate, fumarate, tartrate, citrate, benzoate, 4-methoxybenzoate, 2- or 4-hydroxybenzoate, 4-chlorobenzoate, p-toluenesulphonate, methanesulphonate, ascorbate, salicylate, acetate succinate, lactate, glutarate, gluconate, tricarballylate, hydroxynaphthalenecarboxylate or oleate. Formoterol is preferably used in the form of its fumarate salt and as a dihydrate.

The ratio of formoterol to budesonide used according to the invention is preferably within the range of 1:4 to 1:70. The two drugs may be administered separately in the same ratio.

The intended dose regimen is a twice daily administration, where the suitable daily dose of formoterol is in the range of 6 to 100 µg with a preferred dose of 6-48 µg and the suitable daily dose for budesonide is 50 to 4800 µg with a preferred dose of 100-1600 µg. The particular dose used will strongly depend on the patient (age, weight etc) and the severity of the disease (mild, moderate, severe asthma etc).

For administration, the combination is suitably inhaled from a nebulizer, from a pressurized metered dose inhaler or as a dry powder from a dry powder inhaler (e.g. as sold under the trade mark Turbuhaler) or from a dry powder inhaler utilizing gelatine, plastic or other capsules, cartridges or blister packs.

A diluent or carrier, generally non-toxic and chemically inert to the medicament e.g. lactose, dextran, mannitol or glucose or any additives that will give the medicament a desired taste, can be added to the powdered medicament.

Examples of the preparation of suitable dosage forms according to the invention include the following: Formoterol fumarate dihydrate and budesonide (optionally premicronized)

23940-804(S)

7

are mixed in the proportions given above. The agglomerated, free-flowing micronized mixture may be filled into dry powder inhaler such as sold under the trade mark Turbuhaler. When a capsule system is issued, it is desirable to include a filler  
 5 in the mixture.

The micronized mixture may be suspended or dissolved in a liquid propellant mixture which is kept in a container that is sealed with a metering valve and fitted into a plastic actuator. The propellants used may be chlorofluorocarbons of  
 10 different chemical formulae. The most frequently used chlorofluorocarbon propellants are trichloromonofluoromethane (propellant 11), dichlorodifluoromethane (propellant 12), dichlorotetrafluoroethane (propellant 114), tetrafluoroethane (propellant 134a) and 1,1-difluoroethane (propellant 152a).  
 15 Low concentrations of a surfactant such as sorbitan trioleate, lecithin, disodium dioctylsulphosuccinate or oleic acid may also be used to improve the physical stability.

The invention is further illustrated by way of example with reference to the following Examples.

20 Example 1 - Dry powder inhaler (Turbuhaler)

<u>Active ingredient</u>	<u>Per dose</u>
Formoterol (as fumarate dihydrate)	12 µg
Budesonide	200 µg

The storage unit of the inhaler is filled with sufficient for  
 25 at least 200 doses.

23940-804(S)

8

<u>Active ingredient</u>	<u>Per dose</u>
Formoterol (as fumarate dihydrate)	24 µg
Budesonide	200 µg

- 5 The storage unit is filled with sufficient for at least 200 doses.

<u>Active ingredient</u>	<u>Per dose</u>
Formoterol (as fumarate dihydrate)	12 µg
Budesonide	100 µg

- 10 The storage unit is filled with sufficient for at least 200 doses.

Example 2 - Metered dose inhaler

<u>Active ingredient</u>	<u>Per dose</u>
Formoterol (as fumarate dihydrate)	12 µg
15 Budesonide	200 µg
Stabilizer	0.1 - 0.7 mg
Propellant	25 - 100 µg

<u>Active ingredient</u>	<u>Per dose</u>
Formoterol (as fumarate dihydrate)	24 µg
20 Budesonide	200 µg
Stabilizer	0.1 - 0.7 mg
Propellant	25 - 100 µg

<u>Active ingredient</u>	<u>Per dose</u>
Formoterol (as fumarate dihydrate)	12 µg
25 Budesonide	100 µg
Stabilizer	0.1 - 0.7 mg
Propellant	25 - 100 µg

23940-804 (S)

Example 3 - Metered dose dry powder formulation

<u>Active ingredient</u>		<u>Per dose</u>
Formoterol (as fumarate dihydrate)		12 µg
5	Budesonide	200 µg
	Lactose	up to 5, 12.5 or 25 mg
<u>Active ingredient</u>		<u>Per dose</u>
Formoterol (as fumarate dihydrate)		24 µg
Budesonide		200 µg
10	Lactose	up to 5, 12.5 or 25 mg
<u>Active ingredient</u>		<u>Per dose</u>
Formoterol (as fumarate dihydrate)		12 µg
Budesonide		100 µg
Lactose		up to 5, 12.5 or 25 mg

23940-804(S)

10

CLAIMS:

1.           A medicament comprising together  
  
              (i) formoterol or a physiologically acceptable salt thereof, or a solvate of said salt, or a solvate of formoterol;  
5   and  
  
              (ii) budesonide,  
  
for simultaneous administration by inhalation.
2.           A medicament according to claim 1, which comprises, as component (i), formoterol fumarate dihydrate.
- 10 3.           A medicament according to claim 1 or 2, which comprises a non-toxic diluent or carrier.
4.           A medicament according to claim 3, in which the non-toxic diluent or carrier is lactose.
5.           A medicament according to any one of claims 1 to 4,  
15 in which components (i) and (ii) are in dry powder form.
6.           A medicament according to any one of claims 1 to 5, in which the ratio of the component (i) to component (ii) is in a range of 1:4 to 1:70.
7.           A medicament according to any one of claims 1 to 6,  
20 which contains component (i) and (ii) in unit dose form.
8.           A medicament according to any one of claims 1 to 7, in which components (i) and (ii) are for administration by inhalation in the treatment of a respiratory disorder.
9.           A medicament according to any one of claims 1 to 8,  
25 in which components (i) and (ii) are for administration by inhalation in the treatment of asthma.

23940-804(S)

11

10. A medicament according to any one of claims 1 to 9, in which the administration is by inhalation from a dry powder inhaler.

11. A medicament according to any one of claims 1 to 9,  
5 in which the administration is by inhalation from a pressurized metered dose inhaler.

12. A medicament according to any one of claims 1 to 9, in which the administration is by inhalation from a nebulizer.

13. A pharmaceutical composition for administration by  
10 inhalation, which comprises, together for simultaneous administration:

(i) formoterol or a physiologically acceptable salt thereof, or a solvate of the salt, or a solvate of formoterol; and

15 (ii) budesonide.

14. A composition according to claim 13 which comprises, as component (i), formoterol fumarate dihydrate.

15. A composition according to claim 13 or 14 which comprises a non-toxic diluent or carrier.

20 16. A composition according to claim 15 in which the non-toxic diluent or carrier is lactose.

17. A composition according to any one of claims 13 to 16 in which the ratio of the component (i) to component (ii) is in a range of 1:4 to 1:70.

25 18. A composition according to any one of claims 13 to 17 in unit dose form.

23940-804(S)

12

19. A composition according to any one of claims 13 to 18 in which components (i) and (ii) are each in micronised form.

20. A composition according to claim 19 which is in agglomerated form.

5 21. A dry powder inhaler which contains a composition as claimed in claim 20.

22. A capsule, cartridge or blister pack for a dry powder inhaler, the capsule, cartridge or blister pack containing a composition as claimed in claim 20.

10 23. A composition as claimed in any one of claims 13 to 20 for use in the treatment by inhalation of a respiratory disorder.

24. A composition as claimed in any one of claims 13 to 20 for use in the treatment by inhalation of asthma.

15 25. Use, simultaneously, of effective amounts of

(i) formoterol, or a physiologically acceptable salt thereof, or a solvate of said salt, or a solvate of formoterol; and

(ii) budesonide,

20 for treating a respiratory disorder.

26. A use according to claim 25, in which component (i) is formoterol fumarate dihydrate.

27. A use according to claim 25 or 26 in which the ratio of the component (i) to component (ii) is in a range of 1:4 to  
25 1:70.

28. A use according to any one of claims 25 to 27 in a dose regime, wherein the dose regime is twice daily and

23940-804(S)

13

component (i) is used in a dose of 6 - 100  $\mu\text{g}$  per day and  
component (ii) is used in a dose of 50 - 4800  $\mu\text{g}$  per day.

29. A use according to claim 28 in which the dose of  
component (i) is 6 - 48  $\mu\text{g}$  per day and the dose of component  
5 (ii) is 100 to 1600  $\mu\text{g}$  per day.

30. Use of effective amounts of

(i) formoterol, or a physiological acceptable salt  
thereof, or a solvate of said salt, or a solvate of formoterol;  
and

10 (ii) budesonide,

for manufacturing a medicament for treating a respiratory  
disorder, wherein components (i) and (ii) are to be used  
simultaneously to treat the respiratory disorder.

31. A use according to claim 30, in which component (i)  
15 is formoterol fumarate dihydrate.

32. A use according to claim 30 or 31 in which the ratio  
of the component (i) to component (ii) is in a range of 1:4 to  
1:70.

33. A use according to any one of claims 30 to 32, in  
20 which the medicament is to be used in a twice daily dose regime  
wherein component (i) is to be used in a dose of 6 - 100  $\mu\text{g}$  per  
day and component (ii) is to be used in a dose of 50 - 4800  $\mu\text{g}$   
per day.

34. A use according to claim 33 in which the dose of  
25 component (i) is 6 - 48  $\mu\text{g}$  per day and the dose of component  
(ii) is 100 to 1600  $\mu\text{g}$  per day.